

Lipopolysaccharide structure impacts the entry kinetics of bacterial outer membrane vesicles into host cells

O'Donoghue, Eloise; Sirisaengtaksin, Natalie; Browning, Douglas; Bielska, Ewa; Hadis, Mohammed; Fernandez-Trillo, Francisco; Alderwick, Luke; Jabbari, Sara; Krachler, Anne Marie

DOI:

[10.1371/journal.ppat.1006760](https://doi.org/10.1371/journal.ppat.1006760)

Document Version

Peer reviewed version

Citation for published version (Harvard):

O'Donoghue, E, Sirisaengtaksin, N, Browning, D, Bielska, E, Hadis, M, Fernandez-Trillo, F, Alderwick, L, Jabbari, S & Krachler, AM 2017, 'Lipopolysaccharide structure impacts the entry kinetics of bacterial outer membrane vesicles into host cells', *PLoS pathogens*, vol. 13, no. 11, pp. e1006760.
<https://doi.org/10.1371/journal.ppat.1006760>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

**Lipopolysaccharide structure impacts the entry kinetics of bacterial outer
membrane vesicles into host cells**

Eloise J O'Donoghue¹, Natalie Sirisaengtaksin², Douglas F. Browning¹, Ewa Bielska¹,
Mohammed Hadis³, Francisco Fernandez-Trillo³, Luke Alderwick¹, Sara Jabbari^{1,4} and Anne
Marie Krachler^{2,*}

¹Institute of Microbiology and Infection, School of Biosciences, University of Birmingham,
Edgbaston, B15 2TT Birmingham, UK

²Department of Microbiology and Molecular Genetics, University of Texas McGovern Medical
School at Houston, Houston, TX, 77030, USA.

³Institute of Microbiology and Infection, School of Chemistry, University of Birmingham,
Edgbaston, B15 2TT Birmingham, UK

⁴School of Mathematics, University of Birmingham, Edgbaston, B15 2TT, Birmingham, UK

*Correspondence to: Anne Marie Krachler (anne.marie.krachler@uth.tmc.edu)

This file contains:

Supporting Materials and Methods

Supporting Figures S1-S6

SUPPORTING MATERIALS AND METHODS

Nanoparticle tracking analysis

After purification, OMV samples were diluted 1×10^{-6} in filtered sterile PBS. Particle diameter and concentration were measured using the Nanosight LM10 particle tracking analysis, with a minimum of 100 tracks per sample, performed in triplicate. Camera shutter 1495 and gain of 450 were used, and size distribution scatter plots were created using GraphPad Prism. Size distribution was analysed using analysis of variance (ANOVA) with a Brown Forsythe test for equal variance.


Measurement of ζ -potential as an indicator of OMV surface charge

700 μ l of OMV preparations were analysed using a Zeta Sizer (Malvern Instruments) and data from an average of 30 readings/sample were acquired at 37 °C and means were plotted.

Visualization of outer membrane vesicles by Transmission Electron Microscopy

10 μ l of isolated outer membrane vesicles in sterile deionized distilled water were added to 400-mesh copper grids, and negatively stained with 4% uranyl acetate for 2 min. Samples were then observed using a Jeol 1200Ex transmission electron microscope (Birmingham Electron Microscopy Facility) with an acceleration of 75kV.

41 SUPPORTING FIGURES



42

43 **Figure S1. Morphology, size, charge and probe orientation of reporter OMVs.** (A) Electron

44 micrographs of negative stained OMV fractions from EHEC wt (left image) or EHEC ClyA-Bla

45 (centre and right images). Scale bars, 0.5 μm . (B) Isolated OMVs were diluted 1×10^{-6} fold and

46 nanoparticle tracking analysis was used to determine the size distribution. Black lines represents

47 median size from at least 200 tracks acquired per sample. Statistical significance was determined

48 by ANOVA, with a Brown Forsythe test to determine equal variance. (***) $p \leq 0.005$, (ns) not

49 significant. (C) ζ -potentials of isolated OMVs. Values represent means from 30 readings per

50 sample. Only means are displayed since individual readings are not accessible instrumentally. (D)

51 OMV fractions from EHEC expressing Cly-Bla, Bla-ClyA or carrying empty vector were treated

52 with papain for 30 or 60 minutes, and used for Western Blotting with α -Bla antibody.




Figure S2. Rates of uptake/dismantling and concentration dependency of uptake kinetics for OMVs. (A) CCF2-AM loaded Hela cells exposed to EHEC OMVs carrying ClyA-Bla (red), or empty vector (grey) at an MOI of 1000 for 3 h. Rate of uptake over time was extracted from data in Figure 2A and data shown are means \pm stdev ($n=3$). (B) FRET change upon exposure of Hela cells to EHEC OMVs carrying ClyA-Bla (reporting on exposure to OMV surface to cytoplasm) or Bla-ClyA (reporting on exposure of luminal cargo to cytoplasm). (C) Hela cells were exposed to EHEC or K12 ClyA-Bla OMVs at an MOI of 1000 for 3 hours. Rates of uptake over time were extracted from data in Figure 3A and are means \pm stdev ($n=3$). (D) Experiments were repeated as above but using different OMV concentrations (0-20 $\mu\text{g/ml}$ of protein, corresponding to an MOI of 0- 2000), and maximum rates (D) and efficiency of uptake (E) determined as described above. Data are means \pm stdev ($n=3$).




Figure S3. Uptake for OMVs from serotypes O157, O42 and O16 with or without O antigen. CCF2-AM loaded RKO intestinal epithelial cells were exposed to OMVs from EHEC O157 (A), EAEC O42 (B), and K12 O16 (C), with O antigen (red) and without O antigen (blue), at an MOI of 1000 for 3 hours. FRET changes (blue/green fluorescence, A-C) and efficiency of uptake (total change over three hours, D) are shown as means \pm stdev (n=3).




Figure S4. Rates of uptake for OMVs from serotypes O157, O42 and O16 with or without O antigen. CCF2-AM loaded Hela cells were exposed to OMVs from EHEC O157 (A), EAEC O42 (B), and K12 O16 (C), with O antigen (red) and without O antigen (blue), at an MOI of 1000 for 3 hours. Polynomials were fitted to each data set using the cubic spline function csaps in Matlab. Numerical estimates of the gradients of the resulting polynomials were determined using the gradient function. Data shown are means \pm stdev (n=3).




Figure S5. Effect of blebbistatin and dynasore on uptake of OMVs. HeLa cells were either left untreated or pre-treated 80 μ M Dynasore for dynamin inhibition (grey), or 20 μ M blebbistatin for macropinocytosis inhibition (orange) for 1h at 37 $^{\circ}$ C and exposed to ClyA-Bla OMVs isolated from EHEC (A, B), EAEC (C, D), or K12 (E, F) at an MOI of 1000 for 3 hours. The FRET signal (ratio of blue:green fluorescence) over time was plotted as mean \pm stdev (n=3).




Figure S6. Effect of pharmacological treatments on OMV uptake. HeLa cells were either left untreated or pre-treated with 5 ug/ml papain (lilac), 1 ug/ml chlorpromazine (pink), 5mM methyl- β -cyclodextrin (light green) or 1 μ g/ml filipin (turquoise) and exposed to ClyA-Bla OMVs isolated from EHEC (A, B), EAEC (C, D), or K12 (E, F) at an MOI of 1000 for 3 hours. The FRET signal (ratio of blue:green fluorescence) over time was plotted as means \pm stdev (n=3).